


E Joffe MD MSc¹, G Nowakowski MD², H Tun MD³, A Rosenthal DO⁴, M Lunning DO⁵, R Ramchandren MD⁶, C Li DDS DMSc⁷, L Zhou PhD⁷, E Martinez PhD⁷, R von Roemeling MD⁷, R Earhart MD PhD⁷, M McMahon⁷, I Isufi MD⁸, and L Leslie MD⁹


¹Lymphoma Service, Memorial Sloan Kettering Cancer Center, New York, NY; ²Division of Hematology, Department of Internal Medicine, Mayo Clinic-Minnesota, Rochester, MN; ³Division of Hematology/Oncology, Department of Internal Medicine, Mayo Clinic-Florida, Jacksonville, FL; ⁴Department of Hematology, Mayo Clinic-Arizona, Phoenix, AZ; ⁵University of Nebraska, Omaha, NE; ⁶University of Tennessee Medical Center, Knoxville, TN; ⁷Curis Inc., Lexington, MA; ⁸Yale New Haven Hospital, New Haven, CT; ⁹John Theurer Cancer Center, Hackensack, NJ

INTRODUCTION

BCR and TLR Pathways independently drive NF-κB overactivity (IMBRUVICA Package Insert, Rev 08/2018)



In preclinical testing, blocking both IRAK4 and BTK drove tumor reduction better than blocking either one alone (Booher et al. Waldenström Roadmap Symposium 2019)



IRAK4 is essential for TLR and IL-1R signaling in B-cell proliferation forming multiprotein complexes causing inflammation and tumor growth (1, 2)


Emavusertib, a novel oral IRAK4 inhibitor, dosed twice daily has previously shown:

- Single agent activity in R/R non-Hodgkin Lymphoma patients (3)
- Demonstrated in preclinical studies to overcome tumor resistance to ibrutinib and PI3K inhibitors (4)
- Crossed the blood-brain barrier, reversed IRAK4 pathway activity and caused tumor regression, including cure in a murine PDX model with transplanted A20 NHL to the brain (5)
- Showed in-vivo synergy in B-cell NHL in combination with ibrutinib (6)

STUDY DESIGN

TakeAim-Lymphoma (NCT03328078)

- Part A1:** dose escalation of emavusertib as monotherapy
- Part A2:** dose escalation of emavusertib in combination with ibrutinib
 - Endpoints of Part A1 and A2 include safety, tolerability, and RP2D
 - Treatment: Oral, once-daily (QD), or twice-daily (BID), dosing in continuous 21-day cycles
- Part B:** expansion cohorts of emavusertib in combination with ibrutinib (not yet initiated)



Abbreviations
Relapsed/Refractory (R/R), Marginal Zone Lymphoma (MZL), Diffuse Large B-Cell Lymphoma (DLBCL), Primary Central Nervous System Lymphoma (PCNSL), Non-Hodgkin Lymphomas (NHL), Mantle Cell Lymphoma (MCL), Complete Response (CR), Partial Response (PR), Stable Disease (SD), Progressive Disease (PD), Recommended Phase II Dose (RP2D)


RESULTS

- 13 patients received emavusertib + ibrutinib combination therapy: 8 patients discontinued treatment due to adverse event (2), PD (3), and other (3)
- Emavusertib in combination with ibrutinib is well tolerated
- No DLTs observed at 200mg, 2 DLTs observed at 300mg (Stomatitis and Syncope)


Combination therapy	Total (N = 13)	Grade 3+ Treatment-Related Adverse Event	Safety			
			Emavusertib 200 mg BID ibrutinib 420 mg QD (N = 1)		Emavusertib 300 mg BID ibrutinib 560 mg QD (N = 4)	
Female n : Male n	6 : 7	# patients having grade 3+ TRAEs	1	4	1	3
Age (yrs): median (range)	66 (56, 92)	Platelet count decreased		2	1	
Diagnosis		Alanine aminotransferase increased	1			
CLL	2	Anaemia		1		
PCNSL	2	Blood alkaline phosphatase increased		1		
DLBCL	2	Diarrhea				1
MCL	2	Fatigue		1		
MZL	3	Hyponatraemia	1			
WM	2	Lipase increased		1		
Prior lines of therapy: median (range)	3 (1-8)	Muscular weakness				1
Prior BTK inhibitor	6	Pain				1
		Stomatitis				1
		Syncope			1	
		Thrombocytopenia				1
		Vomiting				1

Evaluable Patients treated with emavusertib + ibrutinib

- 4 patients that received prior BTK treatment show promising anti-cancer activity (SD/CR)
- 4/13 patients were not evaluable for tumor burden; 1 patient progressed without evaluable tumor burden; 3 patients had no response assessments prior to discontinuation of treatment (1 due to adverse event, 1 died, 1 other)




- Majority of patients had decreases in tumor burden or SD over time



RESULTS

Preliminary Efficacy Data From Patients with Combination Therapy



Data extracted May 6th, 2022

SUMMARY

- In prior clinical studies oral emavusertib monotherapy appears to be well tolerated with an acceptable long term safety profile
- Preliminary data suggest that combination therapy may overcome ibrutinib resistance in hematological malignancies
- Objective responses occurred at both 200mg and 300mg BID dose levels. All responding patients are presently being treated at 200mg BID of emavusertib with full dose of ibrutinib

ACKNOWLEDGEMENTS

We would like to thank the patients, their families and caregivers for their invaluable contribution and participation in this study.

REFERENCES

1. Küppers et al. J Exp Med. 2015;212 (13):2184
2. Smith et al. Nat Cell Biol. 2019;21 (5):640-50
3. Nowakowski et al. Blood. 2020;36 (Suppl 1):44-45
4. Guidetti et al. AACR Mol Cancer Ther. 2021;20 (Suppl 12):P073
5. Von Roemeling et al. AACR; Mol Cancer Ther. 2021;20 (Suppl 12):P243
6. Booher et al. Waldenström Roadmap Symposium. 2019

CONTACT INFORMATION

Reinhard von Roemeling, MD
Senior VP, Clinical Development
rvonroemeling@curis.com

Curis, Inc.
128 Spring Str
Building C – Suite 500
Lexington, MA 02421
www.curis.com

